


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : D06P 3/26, 3/54, 3/72, 3/79, 3/82, C09B 62/80, 1/20, 1/50, 1/56, C08K 5/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 95/21958 (43) International Publication Date: 17 August 1995 (17.08.95)</p>
<p>(21) International Application Number: PCT/GB95/00097 (22) International Filing Date: 23 January 1995 (23.01.95) (30) Priority Data: 9402607.7 10 February 1994 (10.02.94) GB (71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB). (72) Inventor; and (75) Inventor/Applicant (for US only): HALL, Nigel [GB/GB]; 35 Newton Drive, Greenmount, Bury BL8 4DH (GB). (74) Agents: GILES, David, Eric et al.; Intellectual Property Group, Zeneca Specialties, P.O. Box 42, Hexagon House, Blackley, Manchester M9 8ZS (GB).</p>		<p>(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ). Published With international search report.</p>
<p>(54) Title: ANTHRAQUINONE DYES CONTAINING A FLUOROSULPHONYL GROUP AND USE THEREOF</p> <p>(57) Abstract</p> <p>A process for colouring a synthetic textile material or fibre blend thereof which comprises applying to the synthetic textile material a compound of formula (1) wherein Ring A and Ring D are optionally substituted and at least one of Ring A or Ring D carries at least one -SO₂F group except for 2-[(9,10-dihydro-9,10-dioxo-1-anthracenyl)amino]-ethanesulphonylfluoride, 2-[(9,10-dihydro-9,10-dioxo-2-anthracenyl)amino]-ethanesulphonylfluoride, 2,2'-[(9,10-dihydro-9,10-dioxo-1,4-anthracenediyl)diimino]bisethanesulphonylfluoride, 2,2'-[(9,10-dihydro-9,10-dioxo-1,5-anthracenediyl)diimino]bisethanesulphonylfluoride, 2,2'-[(9,10-dihydro-2-methoxy-9,10-dioxo-1,4-anthracenediyl)diimino] bisethanesulphonylfluoride, 2-[(4-(benzoylamino)-9,10-dihydro-9,10-dioxo-1-anthracenyl)amino] ethanesulphonylfluoride, 2-[(9,10-dihydro-4-(methylamino)-9,10-dioxo-1-anthracenyl)amino]ethane sulphonylfluoride.</p> <div style="text-align: right;">  <p>(1)</p> </div>		

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Anthraquinone dyes containing a fluorosulphonyl group and use thereof

The present invention relates to a process for colouring synthetic textile materials, to synthetic textiles when coloured, to a process for the mass coloration of plastics, to plastics when coloured to certain novel anthraquinone dyes and to compositions containing anthraquinone dyes.

According to the present invention there is provided a process for colouring a synthetic textile material or fibre blend thereof which comprises applying to the synthetic textile material a compound of Formula (1):



Formula (1)

wherein:

Ring A and Ring D are optionally substituted and at least one of Ring A or Ring D carries at least one $-SO_2F$ group except for

- 2-[(9,10-dihydro-9,10-dioxo-1-anthracenyl)amino]-ethanesulphonylfluoride,
- 2-[(9,10-dihydro-9,10-dioxo-2-anthracenyl)amino]-ethanesulphonylfluoride,
- 2,2'-[(9,10-dihydro-9,10-dioxo-1,4-anthracenediyl)diimino]bisethane sulphonylfluoride,
- 2,2'-[(9,10-dihydro-9,10-dioxo-1,5-anthracenediyl)diimino]bisethane sulphonylfluoride,
- 2,2'-[(9,10-dihydro-2-methoxy-9,10-dioxo-1,4-anthracenediyl)diimino]bisethanesulphonylfluoride,
- 2-[(4-(benzoylamino)-9,10-dihydro-9,10-dioxo-1-anthracenyl)amino]ethanesulphonylfluoride,
- 2-[(9,10-dihydro-4-(methylamino)-9,10-dioxo-1-anthracenyl)amino]ethane sulphonylfluoride.

Different compounds of Formula (1) may be mixed or the compounds of Formula (1) may be mixed with dyes which do not contain an $-SO_2F$ group, such mixtures are a feature of the present invention.

The mixtures may be simple physical mixtures or may be mixed crystals formed for example by co-crystallisation. Such mixtures generally show improvement in dyeing properties. Crystalline modifications of compounds of Formula (1) exist and it is intended that the present definition includes such crystalline modifications which may for example be formed by heat treatment.

According to a further feature of the present invention there is provided a process for colouring a polyester textile material or fibre blend thereof which comprises applying to the polyester textile material

2-[(9,10-dihydro-9,10-dioxo-1-anthracenyl)amino]-ethanesulphonylfluoride,
 2-[(9,10-dihydro-9,10-dioxo-2-anthracenyl)amino]-ethanesulphonylfluoride,
 2,2'-[(9,10-dihydro-9,10-dioxo-1,4-anthracenediyl)diimino]bisethane
 sulphonylfluoride,
 2,2'-[(9,10-dihydro-9,10-dioxo-1,5-anthracenediyl)diimino]bisethane
 sulphonylfluoride,
 2,2'-[(9,10-dihydro-2-methoxy-9,10-dioxo-1,4-anthracenediyl)diimino]
 bisethanesulphonylfluoride,
 2-[(4-(benzoylamino)-9,10-dihydro-9,10-dioxo-1-anthracenyl)amino]
 ethanesulphonylfluoride,
 2-[(9,10-dihydro-4-(methylamino)-9,10-dioxo-1-anthracenyl)amino]ethane
 sulphonylfluoride.

The presence of one or more $-SO_2F$ groups in a dye molecule generally improves the properties of that dye and confers surprisingly good wet-fastness and light-fastness properties.

The synthetic textile material may be selected from secondary cellulose acetate, cellulose triacetate, polyamide, polyacrylonitrile and aromatic polyester. The synthetic textile material is preferably polyamide or aromatic polyester, such as polyhexamethylene adipamide or polyethylene terephthalate more preferably aromatic polyester and especially polyethylene terephthalate. Fibre blends may comprise mixtures of different synthetic textile materials or mixtures of synthetic and natural textile materials. Preferred fibre blends are those of polyester-cellulose such as polyester-cotton. The textile materials or blends thereof may be in the form of filaments, loose fibres, yarn, woven or knitted fibres.

The dyes of Formula (1) preferably have low solubility in water, typically less than 1% preferably less than 0.5% and especially less than 0.2% solubility in water. The dyes of Formula (1) are thus free from water solubilising groups such as $-SO_3H$, $-CO_2H$, $-PO_3H$ and quaternary amino.

The dyes of Formula (1), optionally in conjunction with other disperse dyes may be applied to the synthetic textile materials or fibre blends thereof by methods which are conventionally employed in dyeing disperse dyes to such materials and fibre blends. For example, the dyes of Formula (1) in the form of an aqueous dispersion may be applied by dyeing, padding or printing processes using the conditions and additives conventionally used in carrying out such processes.

The process conditions may be selected from the following:

- i) exhaust dyeing at a pH of from 4 to 6.5, at a temperature of from 125°C to 140°C for from 10 to 120 minutes and under a pressure of from 1 to 2 bar, a sequestrant may be optionally be added;

- ii) continuous dyeing at a pH of from 4 to 6.5, at a temperature of from 190°C to 225°C for from 15 seconds to 5 minutes, a migration inhibitor may optionally be added;
- iii) printing direct at a pH of from 4 to 6.5, at a temperature of from 160°C to 185°C for 4 to 15 minutes for high temperature steaming, or at a temperature of from 190°C to 225°C for 15 seconds to 5 minutes for bake fixation with dry heat or at a temperature of from 120°C to 140°C and 1 to 2 bar for 10 to 45 minutes for pressure steaming, wetting agents and thickeners (such as alginates) of from 5 to 100% by weight of the dye may be optionally be added;
- iv) discharge printing (by padding the dye onto the textile material, drying and overprinting) at a pH of from 4 to 6.5, migration inhibitors and thickeners may optionally be added;
- v) carrier dyeing at a pH of from 4 to 6.5, at a temperature of from 95°C to 100°C using a carrier such as methylnaphthalene, diphenylamine or 2-phenylphenol, sequesterants may optionally be added; and
- vi) atmospheric dyeing of acetate, triacetate and nylon at a pH of from 4 to 6.5, at a temperature of 85°C for acetate or at a temperature of 90°C for triacetate and nylon for from 15 to 90 minutes, sequesterants may optionally be added.

In all the above processes the compound of Formula (1) is applied as a dispersion comprising from 0.001% to 4% of the compound in aqueous medium.

The present compounds generally provide coloured textile material which shows good fastness to washing, light and heat.

Examples of suitable substituents for Rings A and D are cyano, hydroxy, nitro, halo such as fluoro, chloro or bromo, fluorosulphonyl, trifluoromethyl or alkyl, alkoxy, aryl, aryloxy, heterocycloalkyl, alkylthio, arylthio, heteroarylthio, $-SO_2aryl$, $-NHCOalkyl$, $-NHCOaryl$, $-NHSO_2alkyl$, $-NHSO_2aryl$, $-NHSO_2arylalkyl$, $-OCOalkyl$, $-OCOaryl$, $-COalkyl$, $-COaryl$, $-COOalkyl$, $-COOaryl$, $alkylOCOaryl$, $-SO_2alkyl$, $-SO_2aryl$, $-SO_2arylOH$, $-NR^1R^2$ or $-SO_2NR^1R^2$ in which R^1 and R^2 each independently is -H, alkyl, aryl, alkylaryl, cycloalkyl and $-COaryl$ or R^1 and R^2 together with the -N atom to which they are attached form a 5- or 6-membered ring such as morpholino or piperidino. Each of the alkyl, alkoxy, aryl, aryloxy, heterocycloalkyl, cycloalkyl parts of the substituents for Rings A and D may be optionally substituted by -OH, -CN, -Cl, -F, -Br, $-SO_2F$, alkyl, alkoxy, $-OCOalkyl$, $-SO_2NHalkylalkoxy$, $-OCOphenyl$,

$-alkylN-CO(CH_3)_2$, or $-alkylN-CO(CH_3)_2$, alkoxyethyl, hydroxyethyl or phenyl. Each alkyl in the above substituents is preferably C_{1-10} -alkyl more preferably C_{1-6} -alkyl, each alkoxy is preferably C_{1-10} -alkoxy more preferably C_{1-6} -alkoxy, each of which may be straight or branched chain

alkyl or alkoxy, each aryl is preferably phenyl or naphthyl each aryloxy is preferably phenoxy or naphthoxy, each heterocycloalkyl is preferably triazolyl, each alkylthio is preferably C₁₋₆-alkylthio, each arylthio is preferably phenylthio, each heteroarylthio is preferably
 5 benzothiazolythio and each alkylaryl is preferably C₁₋₆-alkylphenyl especially benzyl, each -COaryl is preferably -COphenyl and each alkyl, alkoxy, aryl or cycloalkyl may carry an -SO₂F substituent, furthermore two adjacent substituents may be combined to form a 5-membered heterocyclic ring.

10 Preferred substituents for Rings A and D are hydroxy, chloro, bromo, fluorosulphonyl or aryloxy, heterocycloalkyl, heteroarylthio each of which is optionally substituted by -OH, -SO₂F, -OCOalkyl,

-SO₂NHalkylalkoxy, -OCOphenyl, -alkylN-CO(CH₂)_n, -NHSO₂aryl or -SO₂arylOH;
 15 NR¹R² or -SO₂NR¹R² in which R¹ and R² each independently is -H, alkyl or aryl each of which is optionally substituted by -SO₂F, -OH, alkoxy or alkoxyalkyl.


Especially preferred substituents for Rings A and D are
 -OH, (4-SO₂F)phenoxy, -NH₂, (4-OH-3-SO₂F)phenyl, -NH(4-SO₂Fphenyl),
 20 -NH(3-SO₂Fbenzyl), -NH(2,4-diSO₂Fphenyl), -SO₂phenyl, (4-SO₂F)phenylthio, -
 (CH₂)_nOCO(4-SO₂Fphenyl), -CO(4-SO₂Fphenyl), -NHphenyl and
 -NH(3-SO₂C₆H₄OSO₂Hphenyl).

The compounds of Formula (1) preferably carry a total of from
 1 to 3 -SO₂F groups, more preferably 1 or 2 -SO₂F groups and especially 1
 25 -SO₂F groups.

The compounds of Formula (1) except for the compounds listed above and the following compounds

1-N-(4-methyl-3-fluorosulphonylphenyl)amino-4-N-methylaminoanthraquinone,
 1-N-methylamino-4-N-benzylamino-2-fluorosulphonylanthraquinone,
 30 1-amino-4-N-benzylamino-2-fluorosulphonylanthraquinone,
 1-amino-5-fluorosulphonylanthraquinone, and
 1-N((4-fluorosulphonyl)benzyl)amino-4-methylaminoanthraquinone are novel
 and accordingly form a further feature of the present invention.

A preferred sub-group of compounds of Formula (1) are those
 35 of Formula (2):



Formula (2)

in which:

R¹ and R² each independently is -H, or alkyl, alkylaryl, aryl or -COaryl optionally substituted by -SO₂F, -OH, alkoxy or alkylOH, or -SO₂aryl optionally substituted by -SO₂F, -SO₂alkylOSO₂H or alkyl;

R³ is -H, -Cl, -Br, -SO₂F or alkyl, alkylOCOaryl, aryl, cycloalkyl, heterocycloalkyl, alkoxy, aryloxy, alkylthio, arylthio, heteroarylthio or -SO₂aryl each of which is optionally substituted by -OH, alkyl, alkoxy, -OCOalkyl, -CH₂N-CO(CH₂)ₛ, -SO₂NHalkylalkoxy, -SO₂F, -COOalkyl or -OCOaryl;

R⁴ is -H, -SO₂F, -Cl, -Br, -CN or aryl or aryloxy each optionally substituted by -OH, -SO₂F or -OCOalkyl;

or R³ and R⁴ together with the carbon atoms to which they are attached form a 5-membered heterocyclic ring;

Z is -H, -OR¹, SR¹, -NR¹R² or -COaryl;

R⁵ and R⁶ each independently is -H, -NO₂, -OR¹, -NR¹R² or -SO₂F; and

R⁶ and R⁷ each independently is -H, -SO₂F, or halo.

Where any of the groups represented by R¹, R², R³, R⁵ or R⁶ is or contain alkyl or alkoxy substituents these are preferably C₁-₁₀-alkyl or C₁-₁₀-alkoxy, more preferably C₁-₆-alkyl or C₁-₆-alkoxy.

Where any of the groups represented by R¹, R², R³, R⁴, R⁵ or R⁶ is or contain aryl or aryloxy substituents these are preferably phenyl or phenoxy.


Where either of R¹ or R² is alkylaryl it is preferably C₁-₆-alkylaryl, more preferably benzyl.

The heterocycloalkyl group represented by R³ is preferably 1,2,4-triazin-5-yl.

The heteroarylthio group represented by R³ is preferably benzothiazolylthio.

Where R⁶ and R⁷ are halo the halo is preferably -Cl, -Br or -I and more preferably -Cl.

Where R³ and R⁴ together with the carbon atoms to which they are attached form a 5-membered heterocyclic ring this is preferably of the Formula (3):



Formula (3)

in which

R⁹ is alkyl optionally substituted by OR¹ wherein R¹ is as hereinbefore defined.

R⁹ is preferably C₁₋₆-alkyl.

Especially preferred compounds of Formula (2) are those in which R¹, R², R⁴, R⁵, R⁶, R⁷ and R⁸ are all -H, Z is -OH and R³ is (4-fluorosulphonyl)phenoxy and in which R¹, R², R⁵, R⁶, R⁷ and R⁸ are all -H, Z is -NH₂ and R³ and R⁴ are both 4-fluorosulphonyl phenoxy.

Compositions comprising dispersions of the compounds of Formula (1) in which Ring A and Ring D are as hereinbefore defined; and at least one of Rings A or D carries directly at least one -SO₂F group or carries a substituent to which at least one -SO₂F group is attached in aqueous media are novel and form a further feature of the present invention. The compositions typically comprise from 1% to 30% of the compound of Formula (1) and are preferably buffered at a pH from 2 to 7, more preferably at a pH from pH 4 to 6.

These dispersions may further comprise ingredients conventionally used in dyeing applications such as dispersing agents for example lignosulphonates, naphthalene sulphonic acid/formaldehyde condensates or phenol/cresol/sulphanilic acid/formaldehyde condensates, surfactants, wetting agents such as alkyl aryl ethoxylates which may be sulphonated or phosphated, inorganic salts, de-foamers such as mineral oil or nonanol, organic liquids and buffers. Dispersing agents may be present at from 10% to 200% on the weight of the compound of Formula (1). Wetting agents may be used at from 0% to 20% on the weight of the compound of Formula (1). The dispersions may be prepared by bead milling the compound of Formula (1) with glass beads or sand in an aqueous medium.

According to a further feature of the present invention there is provided a process for the mass coloration of plastics which comprises incorporating into a plastics material a compound or mixture thereof which is free from water solubilising groups, of Formula (1) wherein Ring A and Ring D are as hereinbefore defined and at least one of Rings A or D carries directly at least one -SO₂F group or carries a substituent to which at least one -SO₂F group is attached.

The plastics may be selected from polystyrene, acrylics, styrene/acrylonitrile mixtures, acrylonitrile/butadiene/styrene mixtures, polycarbonate, polyether-sulphone, nylons, rigid PVC (uPVC) and polypropylene.

5 The compound may be incorporated by blending with granules or powdered plastics material by, for example, dry tumbling or high-speed mixing followed by injection moulding on a screw machine or by conventional compounding/masterbatching techniques. The present dyes generally dissolve or disperse readily in hot plastics melt and provide
10 bright coloration generally with good clarity and good light fastness.

The plastics materials when coloured with the above dyes form a further feature of the present invention.

The compounds of Formula (1) may be obtained by usual methods for the preparation of anthraquinone compounds such as by oxidation of
15 anthracene or substituted anthracenes with potassium dichromate in sulphuric acid or by reaction of a phthalic anhydride with benzene or a substituted benzene in the presence of aluminium chloride followed by ring closure in hot sulphuric acid.

20 Fluorosulphonyl groups may be introduced into the compounds of Formulae (1) or (2) by methods generally available in the literature. For example reaction of a compound of Formulae (1) or (2) with chlorosulphonic acid optionally in the presence of dimethylformamide and thionylchloride at a temperature of from 30°C to 100°C gives the chlorosulphonyl derivative. The chlorosulphonyl derivative may be
25 reacted in boiling aqueous media with potassium fluoride to give the fluorosulphonyl derivative.

Alternatively a compound of Formulae (1) or (2) may be sulphonated with sulphuric acid or oleum to give the sulphonic acid derivative which may be converted to the chlorosulphonyl derivative by
30 reaction, either of the free acid or an inorganic salt thereof, with thionylchloride optionally in the presence of a chlorophosphorus compound such as phosphorus oxychloride or phosphorus pentachloride in an organic liquid such as an aromatic hydrocarbon at a temperature of from 20°C to 110°C. The chlorosulphonyl derivative may then be converted to the
35 fluorosulphonyl derivative as described above.

The compounds of Formulae (1) and (2) are useful for the coloration of synthetic textile materials particularly polyester textile materials and fibre blends thereof to which they impart colours which have excellent wet and light fastness properties.

40 The compounds of Formulae (1) and (2) are also useful for the mass coloration of plastics as described above and impart bright colours generally with good clarity and lightfastness.

The invention is further illustrated by the following Examples.

Example 1Preparation of 1-amino-4-hydroxy-2-(4-fluorosulphonylphenoxy)anthraquinone


To a mixture of 1-amino-4-hydroxy-2-(4-chlorosulphonylphenoxy)anthraquinone (2 parts) and 1,4-dioxane (25 parts), stirring at ambient temperature, was added a solution of potassium fluoride (0.6 parts) in water (5 parts). The mixture was heated to 60°C and stirred at this temperature for 20 minutes. After cooling to 0-5°C, the product was isolated by filtration, washed with water and dried at 50°C to yield: 1-amino-4-hydroxy-2-(4-fluorosulphonylphenoxy)anthraquinone (1.6 parts). $\lambda_{\text{max}} = 515.0\text{nm}$ (CH_2Cl_2).

Example 2Preparation of 1,5-dihydroxy-2-(4'-hydroxy-3'-fluorosulphonylphenyl)-4,8-diaminoanthraquinone

Chlorosulphonic acid (11.4 parts) was stirred at room temperature while 1,5-dihydroxy-2-(4'-hydroxyphenyl)-4,8-diaminoanthraquinone (0.5 parts) was added slowly over 30 mins. After stirring for 30 minutes thionyl chloride (2.5 parts) was added. The reaction mixture was stirred at 40°C for 3 hours. On cooling the reaction mixture was poured into ice/water and the resulting aqueous suspension was filtered to yield the sulphonyl chloride derivative (0.49 parts).

The sulphonyl chloride derivative (1.6 parts) was added to 1,4-dioxane (94 parts) at room temperature before adding potassium fluoride (5.2 parts) in water (111 parts). After heating at 60°C for 4 hours the mixture was cooled, filtered and dried to yield 1,5-dihydroxy-2-(4'-hydroxy-3'-fluorosulphonylphenyl)-4,8-diaminoanthraquinone (1.02 parts). $\lambda_{\text{max}} = 587\text{nm}, 629\text{nm}$.

Dyes of Formula (2) in which R⁶ and R⁷ are -H were prepared by the procedure of Example 2:



Formula (2)

and these are summarised in Table 1:-

E.G.	R ¹	R ²	R ³	R ⁴	Z	R ⁵	R ⁶	λ_{\max} nm
3	4-SO ₂ Fphenyl	H	H	H	-OH	H	H	514.7
4	4-SO ₂ Fphenyl	H	H	H	-NH(4-SO ₂ Fphenyl)	H	H	401, 570
5	H	H	4-SO ₂ Fphenoxy	4-SO ₂ Fphenoxy	-NH ₂	H	H	564.1, 582.7
6	4-SO ₂ Fphenyl	H	H	H	-OH	H	H	-
7	3-SO ₂ Fbenzyl	H	H	H	-H	H	H	-
8	2,4-DiSO ₂ Fphenyl	H	H	H	-NH(4-SO ₂ Fphenyl)	H	H	542
9	2,4-DiSO ₂ Fphenyl	H	-SO ₂ phenyl	H	-NH ₂	H	H	367, 590
10	4-SO ₂ Fphenyl	H	H	H	-H	H	H	
11	H	H	H	H	4-SO ₂ Fphenylthio	H	H	
12	H	H	SO ₂ Fphenylthio	H	4-SO ₂ Fphenylthio	H	H	
13	H	H	H	H	-OH	H	H	
14	H	H	-(CH ₂) ₆ OCO(4-SO ₂ Fphenyl)	H	-OH	H	H	
15	H	H	-(CH ₂) ₆ OCO(4-SO ₂ Fphenyl)	H	-CO(4-SO ₂ Fphenyl)	H	H	449
16	-CO(4-SO ₂ Fphenyl)	H	-(CH ₂) ₆ OCO(4-SO ₂ Fphenyl)	H	-OH	H	H	
17	2,4-DiSO ₂ Fphenyl	H	H	H	-NH ₂	H	H	372, 451

For examples of dyes made see Table 1.

Example 18

Preparation of 1-amino-2-fluorosulphonyl-4-(N-phenyl)aminoanthraquinone

5 1-Amino-(N-phenyl)aminoanthraquinone-2-sulphonic acid (2.5 parts) was heated in pyridine (100 parts) at reflux for 2 hours. The reaction mixture was filtered and the pyridine removed on a rotary evaporator to yield 2.3 parts of the pyridinium salt.

10 The pyridinium salt (1 part) was heated at 75°C for 45 minutes in POCl₃ (20 parts). On cooling, the reaction mixture was poured into ice/water, filtered and the resulting solid was washed several times with water. The sulphonyl chloride was heated at 100°C for 3 hours with potassium fluoride (3 parts) in 1,4-dioxan (47 parts)/water (22 parts). After cooling the reaction mixture was filtered and the solid obtained
15 washed with water to yield 1-amino-2-fluorosulphonyl-4-(N-phenyl)aminoanthraquinone (0.2 parts).

Example 19:

Preparation of 1-amino-2-fluorosulphonyl-4-(N-(3-sulphonylethylsulphonic acid)phenyl)aminoanthraquinone

20 1-Amino-4-(N-(3-sulphonylethylsulphonic acid)phenyl)aminoanthraquinone-2-sulphonic acid (1.2 parts) was heated to reflux in pyridine (814 parts) for 24 hours. The pyridine was removed by rotary evaporator and the resulting solid washed with petroleum ether (b. pt. 60-80°C) to give the dipyridinium salt which was dissolved in DMF (972 parts) and thionyl chloride (340 parts) was added dropwise. The mixture
25 was heated at 80°C for 4-5 hours, cooled, poured into water and filtered to give the sulphonyl chloride derivative.

30 The sulphonyl chloride derivative (1.76 parts) was heated at 80°C for 18 hours with potassium fluoride (13.5 parts) in 1,4-dioxan (468 parts)/water (555 parts). After cooling, the solvents were removed on a rotary evaporator to yield a solid which was washed with water before being dissolved in acetone and screened through silica. The acetone was removed to yield 1-amino-2-fluorosulphonyl-4-(N-(3-sulphonylethylsulphonic acid)phenyl)aminoanthraquinone λ_{\max} = 512nm.
35

Example 20 and 21:

Preparation of 1-amino-2-fluorosulphonyl-4-(N-(4-fluorosulphonyl)phenyl)aminoanthraquinone and 1-amino-2-fluorosulphonyl-4-(N-(2,4-difluorosulphonyl)phenyl)aminoanthraquinone

40 1-Amino-2-chlorosulphonyl-4-(N-phenyl)aminoanthraquinone (1 part) was added to a stirred mixture of chlorosulphonic acid (35 parts) and thionyl chloride (8 parts), and heated at 65°C for 2.5 hours. After cooling to room temperature the reaction mixture was poured into
45 ice/water and filtered to yield a solid which was washed several times

with water. The solid was heated at 100°C for 2 hours with potassium fluoride (40 parts) in 1,4-dioxan (480 parts)/water (440 parts). On cooling, the reaction mixture was filtered and the solid obtained washed with water to give 1-amino-2-fluorosulphonyl-4-(N-(4-fluorosulphonyl)phenyl)aminoanthraquinone and 1-amino-2-fluorosulphonyl-4-(N-(2,4-difluorosulphonyl)phenyl)aminoanthraquinone $\lambda_{\text{max}} = 534\text{nm}$. The compounds may be separated chromatographically.

Example 22 and 23

Preparation of 1,5-dihydroxy-2,6-difluorosulphonyl-4,8-diaminoanthraquinone and 1,5-dihydroxy-2-fluorosulphonyl-4,8-diamino-6-chloroanthraquinone.

1,5-Diamino-3,7-disulphoanthrarufin (2.7 parts) was heated in pyridine (10 parts) at reflux for 2 hours. The solution was then filtered and the pyridine removed on a rotary evaporator to yield the dipyridinium salt (2.5 parts).

The pyridinium salt (0.47 parts) was heated at 75°C for 9 hours in POCl₃ (107 parts). After adding toluene (235 parts) the reaction mixture was heated at 120°C for 3 hours. After cooling, the reaction mixture was filtered and the resulting solid sulphonyl chloride was washed several times with water. The sulphonyl chloride was heated at 100°C for 3 hours with potassium fluoride (1.7 parts) in 1,4-dioxan (59 parts). After cooling, the reaction mixture was filtered and the solid obtained washed with water to give 1,5-dihydroxy-2,6-difluorosulphonyl-4,8-diaminoanthraquinone and 1,5-dihydroxy-2-fluorosulphonyl-4,8-diamino-6-chloroanthraquinone. The product was a mixture of 22 and 23, with 23 being the major component of the mixture.

Example 24:

Preparation of 1-amino-4-(N-(4-fluorosulphonyl)phenyl)aminoanthraquinone

1-Amino-4-bromoanthraquinone (3.3 parts), sodium acetate (5 parts), cupric acetate (0.5 parts) and sulphanilyl fluoride (4 parts) were stirred in o-dichlorobenzene (88 parts) at 190°C for 30 hours. The solution was allowed to cool and filtered, the solid was washed with water several times to give 1-amino-4-(N-(4-fluorosulphonyl)phenyl)aminoanthraquinone.


Example 25

Preparation of 1-amino-4-hydroxy-2-(4-fluorosulphonylphenoxy)anthraquinone

A mixture of 1-amino-4-hydroxy-2-phenoxyanthraquinone (0.2g) was stirred overnight with fluorosulphonic acid (10cm³) and poured onto ice. The precipitated solid was collected by filtration and washed with water and dried to give 1-amino-4-hydroxy-2-phenoxy anthraquinone (0.17g, 70%).

CLAIMS

1. A process for colouring a synthetic textile material or fibre blend thereof which comprises applying to the synthetic textile material a compound of Formula (1):



Formula (1)

wherein:


Ring A and Ring D are optionally substituted and at least one of Ring A or Ring D carries at least one -SO₂F group except for

2-[(9,10-dihydro-9,10-dioxo-1-anthracenyl)amino]-ethanesulphonylfluoride,
 2-[(9,10-dihydro-9,10-dioxo-2-anthracenyl)amino]-ethanesulphonylfluoride,
 2,2'-[(9,10-dihydro-9,10-dioxo-1,4-anthracenediyl)diimino]bisethane
 sulphonylfluoride,
 2,2'-[(9,10-dihydro-9,10-dioxo-1,5-anthracenediyl)diimino]bisethane
 sulphonylfluoride,
 2,2'-[(9,10-dihydro-2-methoxy-9,10-dioxo-1,4-anthracenediyl)diimino]
 bisethanesulphonylfluoride,
 2-[(4-(benzoylamino)-9,10-dihydro-9,10-dioxo-1-anthracenyl)amino]
 ethanesulphonylfluoride,
 2-[(9,10-dihydro-4-(methylamino)-9,10-dioxo-1-anthracenyl)amino]ethane
 sulphonylfluoride.

2. A process for colouring a polyester textile material or fibre blend thereof which comprises applying to the polyester textile material

2-[(9,10-dihydro-9,10-dioxo-1-anthracenyl)amino]-ethanesulphonylfluoride,
 2-[(9,10-dihydro-9,10-dioxo-2-anthracenyl)amino]-ethanesulphonylfluoride,
 2,2'-[(9,10-dihydro-9,10-dioxo-1,4-anthracenediyl)diimino]bisethane
 sulphonylfluoride,
 2,2'-[(9,10-dihydro-9,10-dioxo-1,5-anthracenediyl)diimino]bisethane
 sulphonylfluoride,
 2,2'-[(9,10-dihydro-2-methoxy-9,10-dioxo-1,4-anthracenediyl)diimino]
 bisethanesulphonylfluoride,
 2-[(4-(benzoylamino)-9,10-dihydro-9,10-dioxo-1-anthracenyl)amino]
 ethanesulphonylfluoride,
 2-[(9,10-dihydro-4-(methylamino)-9,10-dioxo-1-anthracenyl)amino]ethane
 sulphonylfluoride.

3. A compound of Formula (1):



Formula (1)

wherein:

Ring A and Ring D are optionally substituted and at least one of Ring A or Ring D carries at least one $-SO_2F$ group except for

2-[(9,10-dihydro-9,10-dioxo-1-anthracenyl)amino]-ethanesulphonylfluoride,

2-[(9,10-dihydro-9,10-dioxo-2-anthracenyl)amino]-ethanesulphonylfluoride,

2,2'-[(9,10-dihydro-9,10-dioxo-1,4-anthracenediyl)diimino]bisethane sulphonylfluoride,

2,2'-[(9,10-dihydro-9,10-dioxo-1,5-anthracenediyl)diimino]bisethane sulphonylfluoride,

2,2'-[(9,10-dihydro-2-methoxy-9,10-dioxo-1,4-anthracenediyl)diimino] bisethanesulphonylfluoride,

2-[(4-(benzoylamino)-9,10-dihydro-9,10-dioxo-1-anthracenyl)amino] ethanesulphonylfluoride,

2-[(9,10-dihydro-4-(methylamino)-9,10-dioxo-1-anthracenyl)amino] ethane sulphonylfluoride,

1-N-(4-methyl-3-fluorosulphonylphenyl)amino-4-N-methylaminoanthraquinone,


1-N-methylamino-4-N-benzylamino-2-fluorosulphonylanthraquinone,

1-amino-4-N-benzylamino-2-fluorosulphonylanthraquinone,

1-amino-5-fluorosulphonylanthraquinone, and

1-N((4-fluorosulphonyl)benzyl)amino-4-methylaminoanthraquinone.

4. A compound of Formula (2):



Formula (2)

in which:

R^1 and R^2 each independently is $-H$, or alkyl, alkylaryl, aryl or $-CO$ aryl optionally substituted by $-SO_2F$, $-OH$, alkoxy or alkylOH, or $-SO_2$ aryl optionally substituted by $-SO_2F$,

-SO₂alkylOSO₂H or alkyl;

R¹ is -H, -Cl, -Br, -SO₂F or alkyl, alkylOCOaryl, aryl, cycloalkyl, heterocycloalkyl, alkoxy, aryloxy, alkylthio, arylthio, heteroarylthio or -SO₂aryl each of which is optionally substituted by -OH, alkyl, alkoxy, -OCOalkyl, -CH₂N-CO(CH₂)₅, -SO₂NHalkylalkoxy, -SO₂F, -COOalkyl or -OCOaryl;

R² is -H, -SO₂F, -Cl, -Br, -CN or aryl or aryloxy each optionally substituted by -OH, -SO₂F or -OCOalkyl;

or R³ and R⁴ together with the carbon atoms to which they are attached form a 5-membered heterocyclic ring;

Z is -H, -OR¹, SR¹, -NR¹R² or -COaryl;

R⁵ and R⁶ each independently is -H, -NO₂, -OR¹, -NR¹R² or -SO₂F; and

R⁶ and R⁷ each independently is -H, -SO₂F or halo;

provided that when R¹ is -H, R² is not -C₆H₄SO₂F and except for 1-N-(4-methyl-3-fluorosulphonylphenyl)amino-4-N-methylaminoanthraquinone,

1-N-methylamino-4-N-benzylamino-2-fluorosulphonylanthraquinone,

1-amino-4-N-benzylamino-2-fluorosulphonylanthraquinone,

1-amino-5-fluorosulphonylanthraquinone, and

1-N((4-fluorosulphonyl)benzyl)amino-4-methylaminoanthraquinone.

5. A process for the mass coloration of plastics which comprises incorporating into a plastics material a compound or mixture thereof which is free from water solubilising groups, of Formula (1) wherein Ring A and Ring D are as hereinbefore defined and at least one of Rings A or D carries directly at least one -SO₂F group or carries a substituent to which at least one -SO₂F group is attached.

INTERNATIONAL SEARCH REPORT

Intern Application No
PCT/GB 95/00097

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6	D06P3/26	D06P3/54	D06P3/72	D06P3/79	D06P3/82
	C09B62/80	C09B1/20	C09B1/50	C09B1/56	C08K5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 D06P C09B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Section Ch, Week 7922 Derwent Publications Ltd., London, GB; Class A60, AN 79-416048 & JP,A,54 050 681 (MITSUBISHI CHEM IND KK) , 20 April 1979 see abstract see example 20	3,4
X	US,A,3 952 029 (KRUTAK SR JAMES J ET AL) 20 April 1976 see column 7, line 1 - line 2; examples	2
A	see the whole document	1,3-5
A	US,A,2 576 037 (R.P. PARKER ET AL.) 20 November 1951 see column 1, line 22 - column 2, line 28	1

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"A" document member of the same patent family

Date of the actual completion of the international search

17 May 1995

Date of mailing of the international search report

26. 05. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Ginoux, C

INTERNATIONAL SEARCH REPORT

Inter 1 Application No
PCT/GB 95/00097

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-3952029	20-04-76	NONE	
US-A-2576037	20-11-51	NONE	